

Trade name Ohta' s Isan Seichoyaku

Ohta' s Isan Seichoyaku improves intestinal disorders (small intestine and large intestine) and normalizes bowel movements by triple action combining intestinal probiotics, intestinal herbal medicines, stomachic herbal medicines and digestive enzyme.



【Features】

1. Improve intestinal environment

Three kinds of intestinal probiotics (Bifidobacterium, Lactomin, Clostridium Butyricum) increase the good probiotics in the large intestine and the small one. It restores the disordered intestinal flora and improves the intestinal environment.

2. Excellent effects for intestinal regulator

In addition to intestinal probiotics, combining with Geranium Herb Extract and Mallotus Japonicus Extract strengthen an excellent effect for soft stools and constipation.

It regulates intestines and improves bowel movements of the person who has frequently a trouble of soft stools or constipation.

3. Reduce the burden to intestinal

At the same time Powdered Gentian improves function of weaken stomach, Biodiastase1000 promotes digestion of starch and protein and reduces the burden to intestinal.

Ingredients Daily dosage (9 Tablets) Bifidobacterium----- ----30 mg Lactomin----- -----30 mg Clostridium Butyricum-----9 0 mg Geranium Herb Extract-----102 mg (Geranium Herb 1,020 mg) Mallotus Japonicus Extract----- 63 mg (Mallotus Japonicus 504 mg) Powdered Gentian----- 51 mg Biodiastase 1000----- --60 mg Additive: l -menthol, Hydroxypropyl Cellulose, Lactose, Silicic Anhydride, Magnesium Stearate, Cellulose, Powdered Hy drogenated Maltose Starch Syrup	Manufacturer	OHTA' S ISAN CO., LTD.		
	Dosage Form	Tablet		
	Item	160 tablets	370 tablets	
	Suggested retail price (tax-excluded)	JPW1,380	JPW2,680	
	Japan Article Number	4987033703060	4987033703077	
	Expiry Date	Three years		
	Directions	Please take the following amount after meals.		
		Age	Dosage/ time	Times / day
		For adults (15 years old and over)	3 tablets	3 times a day
		8-14 years old	2 tablets	
	5-7 years old	1 tablet		
	Under 5 years old	Not to be taken		
Indications	Soft stools, Constipation, Abdominal bloating, Intestinal regulation (Regulation of stools)			



PRECAUTIONS



To consult

1. The following person should consult a physician or a pharmacist before taking the medicine
 - (1) Those who are under treatment by a physician
 - (2) Those who or whose parents or siblings have ever been attacked by symptoms or allergy.
 - (3) Those who have ever been attacked by symptoms of allergy caused by medicines.
2. Discontinue the medication immediately when the following cases have occurred and consult a physician or a pharmacist, showing this package insert.
 - (1) When the following symptoms have occurred after taking the medication:

Areas concerned	Symptoms
Skin	Rash ▪ redness ▪ itching

- (2) When symptoms have not improved even after medication for approximately two weeks.

OHTA'S ISAN CO., LTD. 3-2, Sengoku 2-chome, Bunkyo-ku, Tokyo, 112-0011, Japan
TEL: 03(3944)1311 FAX: 03(3947)8322