

# Morishita Jintan medicinal supplies

Supporting needs of self-medication for your health.



MORISHITA JINTAN CO., LTD.

## Remedy for stomatitis having superior portability. / Spray-type

### ■ Efficacy

Stomatitis, throat irritation, sore throat,  
Swollen throat, hoarse voice

### ■ Direction for use

To spray Cool throat to an affected part,.

### ■ Active ingredient s (Per 1mL)

Azulenesulfonic acid sodium / 0.2mg

- Additives: Sodium citrate hydrate, xylitol, saccharin sodium hydrate, macrogol, ethanol, flavor



Content : 6ml  
Several times a day

## Dental cream will remove your pain due to stomatitis.

### ■ Efficacy

Stomatitis, angular stomatitis, cavity, tooth root flame, toothache due to gingivitis, pyorrhea

### ■ Direction for use

To apply Dental cream to an affected part with clean finger-tip or cotton.

### ■ Active ingredient s (Per 100g)

Dibucaine hydrochloride / 1.0g

Ethyl aminobenzoic acid / 0.3g

Cetylpyridinium chloride / 0.1g

l-menthol / 0.1g

- Additives: Polysorbate 80, povidone, a gelation hydrocarbon, flavor



Content : 5g  
Several times a day

## Dental cream will remove your pain due to pyorrhea.

### ■ Efficacy

Toothache due to gingivitis, pyorrhea, decayed tooth, tooth root flame, stomatitis, angular stomatitis

### ■ Direction for use

To apply Dental cream (T4g) to an affected part with clean finger-tip or cotton.

### ■ Active ingredient s (Per 100g)

Dibucaine hydrochloride / 1.0g

Ethyl aminobenzoic acid / 0.3g

Cetylpyridinium chloride / 0.1g

l-menthol / 0.1g

- Additives: Polysorbate 80, povidone, a gelation hydrocarbon, flavor



**Content : 4g**  
**Several times a day**

**To cure the swelling of your lips, can be used from child to adult.**

■ **Efficacy**

Cheilitis, angular stomatitis, stomatitis, gingivitis

■ **Direction for use**

To apply Dental pill cream to an affected part with clean finger-tip or cotton.

■ **Active ingredient s** (Per 100g)

Prednisolone / 0.2g

Cetyl pyridinium chloride hydrate / 0.1g

- Additives: Copper chlorophyllin sodium, povidone, liquid paraffin, citric acid hydrate, gelation hydrocarbon, flavor



**Content : 5g**  
**Several times a day**

## For curing sunburn and burn.

### ■ Efficacy

Sunburn, burn, wound, crack, rash, inflammation, hemorrhoids

### ■ Direction for use

To apply Panpas to an affected part with clean finger-tip, or apply Panpas to gauze, then put it on the affected part

### ■ Active ingredient s (Per 100g)

Zinc oxide / 6.0g

Phenol / 0.5g

Salicylic acid / 0.1g

Isopropyl methyl phenol / 0.05g

York lecithin / 0.5g

- Additives: Cholesterol, stearyl alcohol, bee wax, purified lanoline, vaseline, flavor



**Content : 15g**  
**2 - 3 times a day**

**To cure impetigo & suppuration, can be used from child to adult.**

■ **Efficacy**

Purulent skin disease

■ **Direction for use**

To apply Onofe F to an affected part with clean finger-tip, or apply Onofe F to gauze, then put it on the affected part

■ **Active ingredient s** (Per 100g)

Sulfadiazine / 5.0g

Zinc oxide / 5.0g

Allantoin / 0.5g

- Additives: d- borneol, macrogol, lanoline, vaseline



**Content : 7g**  
**Several times a day**

**Sterilize and disinfect, especially for shoe sores.**

■ **Efficacy**

To sterilize and disinfect for wounds such as shoe sore, cuts and scratches

■ **Direction for use**

To apply Dental pill cream to an affected part with clean finger-tip

■ **Active ingredient s (Per 100g)**

Isopropyl methyl phenol / 1.0g

lidocaine / 1.5g

l-menthol / 2.0g

- Additives: Propylene glycol, cetanol, myristate isopropyl, sodium lauryl sulfate, stearate sorbitan, paraben, glycerin stearate, hexyl decanol, dimethylpolysiloxane, stearic acid polyoxyl stearate, lauromacrogol



**Content : 8g**  
**Several times a day**



**Can be used soon, whenever you wounds.**

■ **Efficacy**

To sterilize and disinfect for wounds such as cuts, scratches and shoe sore

■ **Direction for use**

To apply SirenQ to an affected part with clean finger-tip, or apply SirenQ to gauze, then put it on the affected part

■ **Active ingredient s (Per 10ml)**

- Dibucaine hydrochloride / 10mg
- Chlorpheniramine Maleic acid / 20mg
- Naphazoline hydrochloride / 10mg
- Benzethonium chloride / 10mg



**Content : 10ml**  
**Several times a day**

## To cure slight bed sore and rash.

### ■ Efficacy

To sterilize and disinfect for bed sore, rash, inflammation of the skin, pompholyx, shoe sore, burn, scratch, injury etc...

### ■ Direction for use

To apply CofulS to an affected part with clean finger-tip, or apply CofulS to gauze, then put it on the affected part

### ■ Active ingredient s (Per 10ml)

Acrinol / 0.5g

Zinc oxide / 5.0g

- Additives: Vitamin A oil, ergocalciferol, dipalmitic acid pyridoxine, tocopherol, dl-camphor, squalene, Olive oil, white vaseline, cetanol, flavor



**Content : 19g**  
**1 - several times a day**

## For your eyebrow growth promotion.

■ **Efficacy**

To improve loss and thin growth of eyebrow and hair

■ **Direction for use**

To apply Pereus to an affected part

■ **Active ingredient s (Per 10ml)**

Methyltestosterone / 0.5g

- Additives: Glycerin, ethanol, cholesterol, flavor



**Content : 2.5ml**  
**1 - 2 times a day**

## The pharmaceutical products which improve "micturition" and " the sensation of residual urine" for women.

### ■ Efficacy

Micturition and sensation of residual urine

### ■ Direction for use

- Only for women (15 years and over)

\*Cannot be used for male.

- To be taken three times a day  
(at intervals of 4 hours or more)

### ■ Active ingredient s (Per 3 tablets)

Flavoxate hydrochloride / 600mg

- Containing corn starch, hydroxypropyl cellulose, low substituted hydroxypropylcellulose, stearic acid Mg, hypromellose, macrogol 6000, titanium oxide, arnauba wax.



**Content 20 tablets**  
**To be 2 times a day**

**To recover from motion sickness. High absorbency and easy to drink.**

■ **Efficacy**

For dizziness, nausea and headache due to suffering from motion sickness.

■ **Direction for use**

- To drink a Pode before 30 min to prevent motion sickness before getting on the vehicle.
- 15 years and over.
- \*Cannot be used for 7 years or younger.
- Can be taken up to 2 times a day

■ **Active ingredients (Per 100mL)**

Scopolamine hydrobromide hydrate / 0.22mg  
 Citric acid caffeine / 80mg  
 pyridoxine hydrochloride / 20mg

- D-sorbitol, dehydroacetic acid Na, flavor, ethanol, vanillin



**Content : 10mL/bottle**  
**Can be taken 2 times a day**

## To make inconspicuous on the scar of skin by injure.

### ■ Efficacy

To protect scar and to support treatment.

### ■ Direction for use

- Please do not use it for an infected wound.
- When the symptoms develop such as rash, flare and itch, please to consult a doctor.
- Stop using it to the weak skin such as mucous membrane and face.

\* Please ask the person in charge regarding the other directions.

### ■ Active ingredients

**Cover** : Polyurethane, acrylic system adhesive

**Pad** : Polyurethane



### Content :

S/ 50mmx80mmx3, M/ 80mmx100mmx3

L/ 100mmx115mmx3

**To make inconspicuous on the scar of skin by injure.**

## ■ Efficacy

Use for wound treatment or surgical incision site.

## ■ Direction for use

- Please do not use it for an infected wound.
- When the symptoms develop such as rash, flare and itch, please to consult a doctor.
- Stop using it to the weak skin such as mucous membrane and face.

\* Please ask the person in charge regarding the other directions.

## ■ Active ingredient s

Pad : Polyurethane



**Content :**

**M/ 40mmx65mmx3, L/ 65mmx70mmx3**

## Big and useful pad for bleeding and wound with the effusion.

### ■ Efficacy

To protect scar and to support treatment and join the skin of wound.

### ■ Direction for use

- Please do not use it for an infected wound.
- When the symptoms develop such as rash, flare and itch, please to consult a doctor.
- Stop using it to the weak skin such as mucous membrane and face.

\* Please ask the person in charge regarding the other directions.

### ■ Active ingredients

**Cover** : Polyurethane, acrylic system adhesive, nonwoven fabric

**Pad** : Nonwoven fabric



### Content :

S/ 50mmx80mmx3, M/ 80mmx100mmx3

L/ 100mmx115mmx3



**For shoe sore and scratches, which is easy putting on and hardly peeling off.**

■ **Efficacy**

Especially for shoe sore and scratches

■ **Direction for use**

- Put it on heel, side of the foot and between toes where you have a pain due to shoe sore.

■ **Active ingredient s**

**Cover** : Polyurethane, acrylic system adhesive,

**Pad** : Nonwoven fabric, Polyurethane film



**Content : 23mmx72mmx122mm x10sheet**