

Trade Name: Ohta' s Isan (Can)

Ohta' s Isan is a gastrointestinal medicine that combines crude drugs, antacids and digestive enzyme. It has mild but excellent effects for the treatment of various unpleasant symptoms such as overdrinking, heartburn and heaviness on the stomach.

<Features>

- ◎The efficient combination of 7 kinds of stomachic crude drugs strengthens gastrointestinal function, and is effective in heaviness on the stomach, loss of appetite and so on.
- ◎To make good use of crude drugs, Ohta' s Isan is manufactured with processing as little as possible.
- ◎Rapid-acting, continuously-acting antacids control gastric acidity and have effect on heartburn and stomachache.
- ◎The aroma of the pure crude drugs will bring about pleasant and refreshed feeling.



Ingredients One Dose (1.3g)

Cinnamon Bark.....	92 mg
Fennel.....	24 mg
Nutmeg.....	20 mg
Clove.....	12 mg
Citrus Unshiu Peel.....	22 mg
Gentian.....	15 mg
Powdered Picrasma Wood.....	15 mg
Sodium Bicarbonate.....	625 mg
Precipitated Calcium Carbonate.....	133 mg
Magnesium Carbonate.....	26 mg
Synthetic Aluminum Silicate.....	273.4 mg
Biodiastase.....	40 mg
Additive: l-Menthol	

Manufacturer	OHTA' S ISAN CO., LTD.		
Dosage Form	Powder		
Item	75g	140g	210g
Suggested retail price (tax-excluded)	JPY680	JPY1,200	JPY1,680
Japan Article Number	4987033904016	4987033904023	4987033904030
Expiry Date	Four years		

Indications

Overdrinking, Heartburn, Heaviness in the stomach, Overeating, Stomachache, Stomach discomfort, Indigestion, For promoting digestion, Loss of appetite, Gastrasthenia, Hyperacidity, Feeling of stomach and abdominal distention, Nausea (nausea due to hangover and overdrinking), Vomiting, Heaviness in the chest, Belching

Directions

Please take the following amount after or between meals.

Age	One dose	Times / day
For adults (15 years old and over)	1.3g	3 times a day
8-14 years old	0.65g	
Under 8 years old	Not to be taken	

※One level attached spoon is 1.3g.
Please use the bar on the back of the inner lid for leveling.



PRECAUTIONS FOR USE

DON' T DO (Observe instructions, otherwise you could make the situation worse or side effect might become easy to be caused.)

1. The following persons should not take the medications; Those who are getting dialysis
2. Avoid continuous use of the medication.



CONSULT

1. The following persons should consult a physician or pharmacist before taking the medication;
 - (1) Those who are under treatment by a physician
 - (2) Those who have ever been attacked by symptoms of allergy caused by medicines
 - (3) Those who have been diagnosed as suffering the following diseases; Renal disorders, **hypothyrea**

2. Discontinue the medication immediately when the following cases have occurred and consult a physician or pharmacist, showing this Package Insert;

- (1) When the following symptoms have occurred after taking the medication;

Areas concerned	Symptoms
Skin	Rash, redness, itching

- (2) When symptoms have not improved even after medication for approximately two weeks,

OHATA'S ISAN CO., LTD. 3-2, Sengoku 2-chome, Bunkyo-ku, Tokyo 112-0011, Japan

TEL 03(3944)1313 FAX 03(3947)8322