

*Please store this instruction leaflet in a safe place for later reference.
Be sure to read the instructions carefully before use.*

HIYA KIOGAN TOI (SUGAR-COATED)

樋屋奇応丸 糖衣

販売名：小児良薬樋屋奇應丸

“**HIYA KIOGAN TOI**”, containing 4 kinds of highly-selected crude drug, mildly improves various symptoms of children, such as night crying, loss of appetite, etc.

“**HIYA KIOGAN TOI**” is suitable for children who don't like the characteristic smell and taste of crude drug.

CAUTIONS FOR USE

[Consult a physician for the following]

1. The following persons should consult a physician, pharmacist, or registered distributor of this medication before using this medication.

A person with any serious symptoms, such as intense diarrhea, high fever, etc.

2. In the following cases, discontinue use immediately, and consult a physician, pharmacist, or registered distributor of this medication. Take the instruction leaflet with you.

(1)Nervousness of children, night crying, peevishness, convulsion, loss pf appetite, or gastrointestinal invalidism does not improve after having used this medication for a month.

(2)Common cold, fever caused by common cold, diarrhea, indigestion, or vomiting milk does not improve after having used this medication 5 or 6 times.

INDICATIONS

Nervousness of children, night crying, peevishness, convulsion, common cold, fever caused by common cold, diarrhea, indigestion, vomiting milk, loss of appetite, gastrointestinal invalidism

COMPOSITION (in 60 pills)

| Components | Content |
|------------|----------|
| Agarwood | 15.00 mg |
| Musk | 3.15 mg |

| | |
|-----------|----------|
| Ginseng | 42.00 mg |
| Bear bile | 0.90 mg |

Additives: Sucrose, Rice Powder, Kanbaiko, Borneo Camphor, Paraben, Macrogol, Titanium Oxide, Talk, Carnauba Wax

DOSAGE AND ADMINISTRATION

| Age | One dosage | Daily dosage |
|--------------|-------------|--|
| 7~15 years | 15~20 pills | 3 times a day before or between each meal. |
| 3~7 years | 10~15 pills | |
| 1~3 years | 3~10 pills | |
| Under 1 year | 1~3 pills | |

[Cautions]

- (1)Please follow the recommended dosage and administration.
- (2)This medication should be given to children only under adult supervision.

CAUTIONS ON HANDLING AND STORAGE

- (1)Keep all medication out of the reach of children.
- (2)Do not transfer this medicine to a different container. (It may be misused or the quality may deteriorate.)
- (3)Store in a sealed container in a cool, dry place free from direct sunlight.